



bscpp

**The British Society for
Colposcopy and Cervical Pathology**

PRESIDENT Miss D Lyons, Imperial Hospital, London, W2 1NY Tel: 0203 312 1796 E-mail: deirdre.lyons2@nhs.net
PRESIDENT-ELECT Mr N Myerson, Bradford Royal Infirmary, Bradford BD9 6RJ Tel: 01274 364 889 E-Mail: Nicholas.Mverson@bthft.nhs.uk
SECRETARY Ms. Jenni Byrom, Birmingham Women's Hospital, Birmingham, B15 2TG, Tel: 0121 472 1377 E-mail: jbyrom@nhs.net
TREASURER Dr. Julia Palmer, Sheffield Hospitals NHS Foundation, Sheffield S10 2JF Tel: 0114 226 9696 E-Mail: Julia.palmer8@nhs.net
CO-ORDINATOR Miss S Parisi, BSCCP, Birmingham Women's Hospital, Birmingham B15 2TG, Tel: 0121 335 8200 E-mail: sharon.parisi@nhs.net

Charity Commission Reg. No: 1188125

TO WHOM IT MAY CONCERN

May 2024

Dear Sir or Madam

A STATEMENT FROM THE PRESIDENT OF THE BSCCP

The practice of colposcopy has been established over many decades and was introduced into the UK in the early 1970's. The techniques in colposcopy are described in hundreds of publications in text books and scientific papers.

The application of dilute acetic acid, 3% to 5%, is required to identify the abnormal areas on the cervix, as the abnormal cells we detect at colposcopy are otherwise invisible to the naked eye. Application of Lugol's iodine is used to identify the extent of the area on the cervix that might be affected by the abnormal cells and helps plan treatment.

The use of these solutions is part of the standard colposcopic examination described in all text books on the subject. To confirm the presence of abnormal cells a punch biopsy may be performed. Bleeding can occur and this can be stopped using silver nitrate or Monsel's solution (ferric subsulphate). The use of these solutions is part of the standard colposcopic examination described in all textbooks on the subject.

Treatment of any abnormal cells is usually performed as an outpatient procedure. To provide adequate pain relief, local anaesthetic infiltration of the cervix is performed. To minimise the discomfort of introducing the anaesthetic agent, a fine needle attached to a dental syringe is used. This technique and use of a dental syringe is described in all standard colposcopy textbooks. Many types of local anaesthetic are used and are acceptable, some will include a 'vaso-pressin' agent to minimise blood loss. In all cases the anaesthetic comes in a pre-filled cartridge. Agents such as lignocaine (1-2%) with or without adrenaline (1 in 80,000, 1 in 200,000) (Lignospan Special, Rexocaine, Xylocaine), Scandonest (3% mepivacaine), Citanest (3% prilocaine) with or without Octapressin and Septanest (4% Articaine with adrenaline 1 in 100,000 or 1 in 200,000) are recognised as acceptable agents by the colposcopy community and textbooks.

There was unanimous recognition at Executive Committee level and at the Annual General Meeting assembly, that all these agents are integral to the practice of colposcopy. The Colposcopy Clinical Professional Group has also endorsed their use on the basis of the principle of 'custom and practice'. These agents have been in use for five decades in routine diagnostic and therapeutic colposcopy and are established as the agents of choice at this time.

The BSCCP endorses their use unequivocally

Deirdre Lyons
BSCCP President